

10/526768

Fig. 1A

AKAP188**Amino acid sequence****M**ERPAAGEIDANKCDHLSRGEEGTGDLETSPV

GSLADLPFAAVDIQDDCGLPDVPQGNVPQGNPKRSKENRGDRNDHVKRK
KAKKDYQPNYFLSIPITNKKITAGIKVLQNSILRQDNRLTKAMVGDGSFH
ITLLVMQLLNEDEVNIGTDALLELKPFVEEILEGKHLTLFPFHGIGTFQGG
VGFVKLADGDHVSALLEIAETAARTFQEKILAGESRTFKPHLTFMKLSK
APMLWFKGVRKIEPGLYEQFIDHRFGEEILYQIDLC SMLKKKQSNQYHC
ESSIVIGEKRKEPEDAELVRLSKRLVENAVLKAVQQYLEETQNKQPGGE
GNSVKAEEGDRNGDGSNNRK

Nucleotide sequence (SEQ No. 1)**ATGG**AGCGCCCCCGCCGCGGGAGAAATAGATGCCAATAAGTGTGA

TCATTTATCAAGAGGAGAGGAAGGGACGGGGGACCTGGAGACCAGCCCTG
TAGGTTCTCTGGCAGACCTGCCGTTTGCTGCCGTAGACATTCAGATGAC
TGTGGACTCCCTGATGTACCTCAAGGAAATGTACCTCAAGGAAACCCAAA
GAGAAGCAAAGAAAATAGAGGCGACAGGAATGATCACGTGAAGAAGAGGA
AGAAGGCCAAGAAAGATTATCAACCCAACTATTTCTGTCCATTCCAATC
ACCAACAAAAAGATTACAGCTGGAATTAAAGTCTTGCAAAATTCGATACT
GAGACAGGATAATCGATTGACCAAAGCCATGGTCCGCGACGGCTCCTTTC
ACATCACCTTGCTAGTGATGCAGCTATTAAACGAAGATGAAGTAAACATA
GGTACCGACGCGCTTTTGGAAGCTGAAGCCGTTCTGTGAGGAGATCCTTGA
GGGGAAGCATCTGACTTTGCCCTTCACGGGATTGGCACTTTCCAAGGTC
AGGTTGGCTTTGTGAAGCTGGCAGACGGAGATCACGTGAGTGCCTCCTG
GAGATAGCAGAGACTGCAAAAAGGACATTTGAGGAAAAGGCATCCTGGC
TGGAGAAAAGCAGAACTTTTAAGCCTCACCTGACCTTTATGAAGCTGTCCA
AAGCACCAATGCTCTGGAAGAAGGGAGTGAGAAAAATAGAGCCTGGATTG
TATGAGCAATTTATCGACCACAGATTTGGAGAAGAAATACTGTACCAAT
AGATCTCTGCTCCATGCTGAAGAAAAACAGAGCAATGGTTATTACCACT
GCGAGTCTTCGATCGTGATCGGTGAGAAGGACCGAAAGGAGCCTGAGGAT
GCTGAACTGGTCAAGCTCAGTAAGAGGCTGGTGGAGAACCCGTGCTCAA
GGCTGTCCAGCAGTACCTAGAAGAGACACAGAAACAAAAGCAGCCGGGGG
AGGGGAACCTCCGTCAAAGCTGAGGAGGGAGATCGGAATGGCGATGGCAGT
GATAACAACCGGAAGTGA

Best Available Copy

Fig. 2: Interaction of AKAP188-CFP
and RII α -YFP




Fig. 3: *Fluorescent Resonance Energy Transfer (FRET)*
between AKAP188-CFP and RII α -YFP




Fig. 4A




Fig. 4B

Bleaching of acceptor YFP (RII α -YFP) results in an increase of fluorescence emitted by donor CFP (AKAP18 δ -CFP)




Fig. 5A: Inhibition of interaction of
AKAP18 δ -CFP and RII α -YFP by peptide S-Ht31




Fig. 5B: Result of inhibition of interaction between AKAP18δ-CFP and RIIα-YFP by S-Ht31




Fig. 6A: Inhibition of interaction of AKAP188-CFP and RII α -YFP by peptide S-Ht31 – decrease of FRET signal




Fig. 6B: Peptide S-Ht31-P fails to inhibit interaction of
AKAP18 δ -CFP and RII α -YFP
- no decrease of FRET signal





Fig. 6C: Changes in FRET signals (ratio 535/480) of AKAP188-CFP and RII α -YFP in HEK293 cells in the presence of peptides S-Ht31 and S-Ht31-P








Fig. 7

Fig. 8A



B







Fig. 9

10/526768






Fig. 10

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.